

NIH Public Access Author Manuscript

Alcohol. Author manuscript; available in PMC 2008 June 1

Published in final edited form as: *Alcohol.* 2007 June ; 41(4): 263–270.

Rates of Ethanol Metabolism Decrease in Sons of Alcoholics Following a Priming Dose of Ethanol

Blair U. Bradford^{2,4,*}, Jennifer K. Jackson⁴, Linda L. Powell^{1,4}, and James C. Garbutt^{1,3,4}

1Division of Clinical Research Services, Dorothea Dix Hospital, Raleigh, NC, 27603-2176

23207 Michael Hooker Research Center, Department of Environmental Sciences and Engineering, CB#7431, University of North Carolina, Chapel Hill, NC 27599-7431, USA

3Department of Psychiatry, University of North Carolina at Chapel Hill, CB #7160, Chapel Hill, N.C., 27599-7160

4Bowles Center for Alcohol Studies, School of Medicine, University of North Carolina, Chapel Hill, NC, 27599-7178, USA

Abstract

Rapid changes in rates of ethanol metabolism in response to acute ethanol administration have been observed in animals and humans. To examine whether this phenomenon might vary by risk for alcoholism, 23 young men with a positive family history of alcoholism (FHP) were compared to 15 young men without a family history of alcoholism (FHN). Rates of ethanol metabolism were measured in all subjects first after an initial ethanol dose (0.85 g/kg) and then, several hours later, a second dose (0.3 g/kg), and the two rates were compared. The two groups of subjects were similar in their histories of ethanol consumption. FHP subjects demonstrated faster initial rates of ethanol metabolism, 148 ± 36 mg/kg/hr, compared to FHN subjects, 124 ± 18 mg/kg/hr, p=.01. However, FHN subjects increased their rate of metabolism by 10 ± 27 percent compared to a decrease of -15 ± 24 percent in FHP subjects, p =.007. Fifty-two percent of the FHP and none of the FHN subjects exhibited a decline in metabolic rate of 20% or more, p=.0008. Since a significant proportion of FHP subjects exhibited a decrease in the second rate of ethanol metabolism, these preliminary data might help to partly explain why FHP individuals differ in their sensitivity to ethanol and are more likely to develop alcohol dependence.

Keywords

alcohol metabolism; alcoholism; family history

INTRODUCTION

Alcoholism affects on the order of 12% of individuals in the United States over a lifetime (Hasin & Grant, 2004). Over the past 30 years evidence has accumulated to indicate that there is a strong genetic component underlying vulnerability to alcoholism (Cloninger, 1987;Kendler et al., 1992), though its phenotypic expression is incompletely defined. Convincing evidence has been presented for a number of biological differences between subjects considered at high (family history positive, FHP) or low (family history negative, FHN) risk for alcoholism

^{*} Corresponding author. Tel.: +1-919-843-2597; fax: +1-919-966-4711 E-mail address bubradfor@email.unc.edu (Blair Bradford)

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

because of family history. For example, FHP subjects have been shown to have a decreased amplitude of the P300 wave in the evoked cortical potential (Hesselbrock et al., 2001) and a decreased sensitivity to some effects of ethanol (Schuckit, 1994) compared to FHN subjects. It is of interest that ethanol metabolism was reported as similar between the FHP and FHN populations in the latter study. These observations suggest that biological differences exist between FHP and FHN individuals prior to the onset of overt alcoholism. Furthermore, such biological markers may represent phenotypic expression of risk for alcoholism.

It was observed a number of years ago that 2.5 hours after administration of a large dose of ethanol, a rapid increase in rates of oxygen uptake and ethanol metabolism occurred in the isolated perfused rat liver. This observation was defined as the Swift Increase in Alcohol Metabolism (SIAM) (Yuki & Thurman, 1980), and has subsequently been shown to have a heritable basis in rats *in vivo* (Thurman et al., 1980), to exist in inbred strains of mice (Thurman et al., 1983), deer mice (Glassman et al., 1989) and in man (Thurman et al., 1989). Thurman et al demonstrated that 20% of the FHN subjects studied exhibited an increase in their rate of ethanol elimination by at least 40% following an initial dose of ethanol. Under these conditions, SIAM was characterized by an increase in elimination (Widmark's β) without changes in the volume of distribution, indicating a true acceleration in the rate of ethanol elimination (Thurman et al., 1989). Subjects who exhibited the SIAM phenomenon reported few problems with alcohol. This latter observation led us to hypothesize that increase of ethanol metabolism in humans with the SIAM phenotype might lead to diminished intoxication time and thus, perhaps, a lower likelihood of developing dependence and associated psychosocial problems.

In a pilot study, nine FHP subjects were examined for the SIAM phenomenon. These subjects self-reported that their fathers had alcohol related problems. In this small group, rates of ethanol metabolism <u>decreased</u> significantly, an average of 21%, after rechallenge with ethanol following an initial dose (Handler et al., 1990). This suggested that individuals at risk for alcoholism might differ in their pattern of metabolism of ethanol after an initial priming dose of ethanol compared to individuals at low risk for alcoholism.

The purpose of the present study was to evaluate the effect of ethanol administration on ethanol metabolism in a population of FHP and FHN subjects carefully screened for verification of family history of alcoholism and with similar rates of reported ethanol consumption.

MATERIALS AND METHODS

Subject Selection

FHP and FHN subjects were identified using direct mail techniques targeted to students at North Carolina State University, Raleigh, N.C. Packets were sent to randomly selected Caucasian male students between the ages of 18 and 25 consisting of a cover letter, a Schuckit Questionnaire (to gather information on personal and familial history of alcohol and drug use and related problems as well as basic demographic data, (Schuckit, 1994)), a consent form, a receipt for a cash payment of \$5.00 for returning the questionnaire, and a return envelope. Several FHP subjects were also recruited from contacts with alcoholic families.

The Schuckit Questionnaire was used to initially categorize subjects as FHP (father alcoholic as determined by drinking-related problems) or FHN (no 1st/2nd degree relative with alcoholrelated problems). FHP and FHN subjects were selected from the database after eliminating subjects who indicated a maternal history of alcoholism, a personal history of heavy alcohol or drug use accompanied by problems, or significant personal medical or psychiatric problems. FHN subjects were selected to approximate the FHP population on the variables of age, body size, and use of alcohol. Confidentiality was maintained throughout the study. Subjects selected for participation in the study were interviewed by a psychiatrist (JCG) and by a research assistant (LLP) with a focus on two major areas: 1) determining personal history of substance abuse problems, psychiatric problems, and medical problems, and documenting alcohol and drug use and 2) documenting family history of substance abuse and/or psychiatric problems. Research Diagnostic Criteria (Spitzer et al., 1978)) were employed for the diagnosis of substance use disorders and major mental illnesses in the subjects and Family History-RDC Criteria (Andreasen et al., 1977) were used to establish the presence of alcoholism in the father and other relatives and to screen for a family history of major mental illnesses.

Ethanol use was calculated based on subjects' self-report of average number of drinks/session and average number of sessions/month for the previous six months and the product of these numbers was used to establish a drinking index. Cigarette smoking was based on self-report of the average number of cigarettes smoked per month for the previous six months. Marijuana use was based on the estimate of lifetime number of times smoked. A height/weight ratio was calculated in cm/kg.

Twenty-three FHP subjects were selected for study and 15 FHN subjects were chosen based on their similarities to the FHP group on the variables described above. None of the subjects had a personal history of alcoholism, drug dependence or major mental illness. The characteristics of the FHP and FHN subjects are shown in Table 1. There were no significant mean differences between groups in age, body size, ethanol use, cigarette use or lifetime marijuana use. 8/23 FHP subjects reported using cigarettes, though only two reported an average consumption of more than five cigarettes/month; 0/15 FHN subjects smoked cigarettes. 15/23 FHP subjects reported having used marijuana compared to 9/15 FHN subjects.

Experimental Procedures

The study was approved by the Committee for the Protection of the Rights of Human Subjects at the University of North Carolina and a consent form was obtained from each subject. Prior to testing, all subjects were briefed on the experimental design and were instructed not to consume ethanol for three days before study. The experimental design used in this study was quite similar to that first described by Wilson and later utilized by Thurman (Wilson et al., 1984) (Thurman et al., 1989). Subjects arrived at the research unit at 0800 hours after consuming a light (low fat) breakfast. Starting at about 0830 subjects consumed ethanol (0.85 g/kg) as carbonated wine over 20 minutes. Following wine consumption, subjects were asked to rinse their mouths with water for 5 minutes to remove ethanol. Breath samples to measure breath ethanol concentrations [BAC] were then collected every 10 minutes using a Smith and Wesson 900A Breathalyzer to estimate blood ethanol levels until peak blood ethanol levels were attained (around 90 mg/dl) and subsequently at about 20 minute intervals until blood ethanol levels declined by 30%. Rates of ethanol metabolism were calculated based on the linear rate of decrease in breath ethanol concentrations as described previously (Widmark, 1932) A second dose of ethanol (0.3 g/kg) was consumed to elevate blood ethanol concentrations, again to around 90 mg/dl, and breath samples were collected as before. Breath ethanol levels were monitored until they declined below 20 mg/dl, then vital signs were monitored and the subject was released if stable. The first rate of ethanol elimination is identified as E1 and the second rate as E_2 . The change in rate between E_1 and E_2 is defined as $(E_2-E_1)/E_1$. "Area under the Curve" (AUC) of the estimated blood ethanol levels was mathematically calculated to estimate tissue exposure to ethanol from the peak following the second ingestion of ethanol to an estimate of a zero ethanol level.

Statistical Methods

All values are given as the mean \pm standard deviation. Analyses were carried out using Statistical Analysis System (SAS) software. For statistical comparisons between different time

points within groups the Student's t-test was used where significance was set at p<0.01. For comparisons between two groups, Fisher's test was used with dichotomous variables, otherwise Satterthwaite's unequal-variance t-test was used. Three-group comparisons were based on one-way analysis of variance; and adjustment for covariables was carried out using the standard analysis of covariance. Results of these parametric procedures were confirmed by nonparametric analyses (details not reported).

RESULTS

Figure 1 depicts a typical experiment for a subject. Following ingestion of ethanol, breath samples were taken at 20 to 30 minute intervals. Rates of elimination were calculated using the decrease in BAC per unit time extrapolated from the linear portion of the elimination curve as described in Methods. In this example, the first rate of ethanol metabolism, E_1 was 141 mg/kg/hr and the second rate, E_2 , was 94 mg/kg/hr with a decrease in rate of -33% in this subject.

Peak BAC₁ and BAC₂ were similar for FHN and FHP subjects (Table 1). E_1 was lower in FHN subjects compared to FHP subjects, t=-2.67, p=.01, (Table 1 and Figure 2), whereas E_2 was not different between FHN and FHP subjects. The change in rate between E_1 and E_2 was not significant for the FHN group, (t=1.42, p=.18) but there was a significant decrease in the FHP group (t=-3.03, p=.006). Furthermore, the mean change in rate was +10 ± 27 percent in FHN subjects and a <u>decrease</u> of -15 ± 24 percent in the FHP group, between group difference, t= 2.91, p=.007.

A dot plot of the individual changes in rate of ethanol metabolism for FHP and FHN subjects is shown in Figure 3. These data cluster into two distinct groups. Two cutoff points are shown on this figure; one for an <u>increase</u> in rate of 40% or greater (SIAM) and the other for a <u>decrease</u> in rate of 20% or more (which we have defined as <u>Decrease In Ethanol Metabolism</u>, DIEM). Our previous work in humans demonstrated that the prevalence of SIAM, a rate increase of 40% or more, occurred in 20% of the FHN subjects studied (Thurman et al., 1989). In the present study, as shown on Figure 3, SIAM was observed in 4 of 15 subjects or 27% of the FHN subjects, only one (4%), demonstrated SIAM (FHP vs. FHN, Fisher's exact test, p=.07). Conversely, DIEM, or a <u>decrease</u> in rate of ethanol metabolism of more than 20% was observed in 0% of FHN subjects and in 52% of FHP subjects (Fisher's exact test, p=. 0008).

Reported ethanol consumption, E_1 and rate change were examined as factors contributing to peak BAC₁, E_1 , and rate change. There was no evidence that reported ethanol consumption was associated with either E_1 (r=.15, p=.37) or rate change (r=-.03, p=.85). E_1 was, however, significantly correlated with rate change, r= -.56, p=.0003. In other words, the higher a subject's initial rate of ethanol metabolism the more likely he was to have a decrease in rate. Because E_1 differed between FHP and FHN subjects we conducted an analysis of covariance controlling for E_1 and examining for group status and rate change. Using this analysis, group status continued to show evidence as a factor in predicting rate change, p=.06.

Examination of peak BAC₁ and E₁ revealed no correlation between these two in either FHN (r=-.34, p=.21) or FHP (r=.14, p=.54) subjects. To examine whether differences in peak BAC₁ affected group differences in rate change an analysis of covariance was conducted. After matching on peak BAC₁ significant group differences were still present in rate change, p=.03.

Estimated area under the curve (AUC) was calculated from peak BAC₂ to time of estimated zero BAC in FHP and FHN subjects. This was used as an additional way in which to validate the rate of metabolism E_1 and E_2 in the event that the linearity of the ethanol decline did not follow a perfect zero order kinetic. The mean AUC₂ was 14,676 ± 5344 mg/dl min for FHN

subjects and 16,628 \pm 5993 mg/dl min for FHP subjects, t=-1.05, p=.30. FHP subjects were subdivided into groups defined as those that exhibited the DIEM phenomenon (FHP-DIEM) and those that did not (FHP-Non DIEM) and the AUC was compared (Figure 4). FHP-DIEM subjects exhibited increased AUC, 19,792 \pm 6003 mg/dl min, compared to FHP-Non-DIEM subjects, 13,177 \pm 3778 mg/dl min, (t=3.19, p=.005) or to FHN subjects, 14,676 \pm 5344 mg/ dl min (t=2.31, p=.03). The characteristics of FHP-DIEM, FHP-Non-DIEM and FHN subjects are shown in Table 2. FHP-DIEM subjects were similar to the other groups in body size and in self-reported ethanol consumption. FHP-DIEM subjects exhibited higher peak BAC₁ and peak BAC₂, faster initial rates of ethanol metabolism, and slower second rates of ethanol metabolism.

DISCUSSION

The present study demonstrates that individuals at high risk for the development of alcoholism, based on family history, are more likely to exhibit a decrease in their ability to metabolize ethanol following an initial priming dose of ethanol than individuals at low risk for alcoholism. The decrease in rate of alcohol metabolism or DIEM is tentatively defined as a decrease of at least 20% in the rate of ethanol metabolism between an initial rate and a second rate. The finding that DIEM is much more prevalent in FHP subjects than FHN subjects is of interest and may have clinical relevance. Perhaps most intriguing is the possibility that when some FHP subjects exhibit a slow rate of ethanol metabolism this would lead to longer tissue exposure to ethanol compared to FHN subjects-- even when both groups have similar consumption patterns (Table 2). Assuming that increased tissue exposure is a factor in the development of pharmacodynamic tolerance, DIEM could represent a new mechanism to explain why some FHP individuals exhibit higher CNS tolerance or develop CNS tolerance faster than FHN individuals.

This study and a previous paper from our group (Thurman et al., 1989) described two phenomena, in humans, after administration of ethanol: a SIAM phenomenon, more commonly found in the FHN population, a metabolic tolerance, that potentially could have a protective effect against alcohol dependence; and a DIEM phenomenon, more commonly found in the FHP population, that might be associated with increased vulnerability to alcoholism.

Differences in initial rate of metabolism

It is important to note that we found a difference in the <u>initial</u> rate of ethanol metabolism between FHP and FHN subjects, with FHP subjects having a faster rate. In contradiction to our findings, Schuckit reported that FHP and FHN subjects had similar rates of ethanol disappearance using the same dose as the present study resulting in similar peak alcohol concentrations in FHN and FHP subjects (Schuckit, 1984). Lower peak values and rate differences in the present study may be the result of experimental design. It is important to note that our experiment and Schuckit's differed in that our subjects ate a light breakfast prior to drinking alcohol whereas Schuckit's subjects consumed alcohol on an empty stomach (10 hour fast). More recently, Schuckit demonstrated that when FHP and FHN subjects were fed a light standardized breakfast the slope of the decline in ethanol concentration was about 33% faster in FHP vs FHN subjects (Schuckit et al., 1996). These data were not discussed per se in this paper but the differences are consistent with the present study and suggest that food consumption prior to alcohol can affect the absorption and metabolism of alcohol.

It is long known from work in the rat that rates of alcohol metabolism cannot be accelerated after fasting (Yuki & Thurman, 1980); however, Wilson elegantly demonstrated that in human subjects, acute metabolic tolerance to ethanol or AMTE can occur after fasting (Wilson et al., 1984). These data evaluated the role of ethnicity in metabolic tolerance with fasted subjects using very similar experimental protocols as the current study. SIAM occurred in 11 subjects

out of 46 in all subjects. Those subjects of Polynesian and Oriental descent exhibited a greater average increase in rate than Caucasian subjects. Additionally, DIEM was observed in 8% of subjects in Wilson's study; however, the ethnicity of the subjects was unclear from the published data.

Why differences were found in initial metabolic rate between FHP and FHN subjects in the present study remains an unanswered question. Some possible reasons that the FHP subjects had a faster initial metabolic rate of alcohol than the FHN subjects include: 1) differences existed between FHP and FHN subjects in prior ethanol use such that FHP subjects had developed a metabolic tolerance; 2) differences in biochemistry between FHP and FHN subjects in their ability to metabolize alcohol. One could speculate that differences in ethanol consumption between FHP and FHN subjects were not reported. It is possible that the FHP subjects were more likely to underreport their alcohol consumption and that they, in fact, consumed more alcohol than FHN subjects; however, we have no way to assess this factor.

Induction of ADH has been reported in the rat using continuous infusion of an alcohol containing diet in a chronic model (Badger et al., 2000). In humans, hepatic ADH II has been reported to contribute from 15% at 5 mM to 40% at 60 mM ethanol to total alcohol metabolism in humans (Li et al., 1977). These data suggest that this enzyme pathway does not require induction and makes a significant contribution to alcohol metabolism. Accelerated metabolism via ADH II may result in the higher initial rates in FHP than FHN individuals. This explanation would be plausible if there was evidence that this form of ADH was correlated with FHP individuals. Choi et al has demonstrated differences in gene expression of ADH 2 and ALDH 2 in FHP alcoholics, FHN alcoholics and normal subjects of Korean dissent. (Choi et al., 2006). These findings are encouraging suggesting that a genetic link providing differential expression of ADH in FHP and FHN subjects exists.

Contributions by gastric ADH for first-pass elimination of alcohol metabolism is an alternative explanation; however, Levitt reports that the contribution of gastric ADH to total alcohol metabolism in humans is inconsequential (Levitt et al., 1997) More recently, a new study carefully evaluated age, gender and alcohol concentration in human gastric mucosa in over 120 patients and concluded that gastric ADH and ALDH are not likely to participate in changes of first pass metabolism (Lai et al., 2000).

ADH activity is cofactor dependent and mitochondrial activation is key in regulation of the NAD⁺/NADH redox state. Yuki observed a peak of hepatic respiration, only when glycogen reserves were high within a few hours after acute ethanol and a return to baseline rates of respiration by about 5-6 hours in the isolated perfused liver (Yuki and Thurman, 1980), supporting the theory that mitochondrial activation and return to normal levels could cause temporal changes in cofactor supply. Additionally, mitochondrial activation can be partially mimicked by adrenergic hormones. Alcohol has been shown to cause an increase in adrenaline and noradrenaline in rodent models (Yuki et al., 1980). Interestingly, noradrenaline levels are reported to be low in abstaining alcoholics as compared to controls and noradrenaline levels did not change when subjects were given stress tests (Ratsma et al, 2002). These data suggest that some of the FHP sons in this study may have inherited an altered or insufficient adrenergic response resulting in poor mitochondrial activation and this link could play a role in the varying cofactor supply to ADH and altering alcohol metabolism.

Chlorzoxazone was used as a marker of cytochrome P450 2E1 activity in non-alcoholic human subjects who were self reported drinkers or non-drinkers. Metabolism of chlorzoxazone was observed to be less in drinkers (Liangpunsakul S, et al., 2005). While it is well known that cytochrome P450 2E1 is one of the 3 primary pathways of alcohol metabolism, and that induction of this pathway occurs after chronic alcohol drinking, or induction can occur by use

of other drugs (Takahashi et al., 1993). It is important to note that in the current study, all subjects were matched based on the fact they reported similar drinking patterns prior to the study (Table 1) and were given the same dose of ethanol. These facts support the theory that cytochrome P450 2E1 might not be induced in the current experiment or be involved in alteration of rates of alcohol metabolism. However, as noted earlier, one cannot rule out that FHP subjects under-reported their drinking compared to FHN subjects and the current data does not rule out genetic variation in expression of CYP 2E1 in FHP or FHN subjects.

One recent interesting report demonstrated that blood catalase levels are higher in FHP than in FHN individuals (Koechling et al., 1995). These findings were independent of alcohol use suggesting that in addition to the other biochemical pathways for elimination of ethanol, catalase may contribute to decrease the blood levels to a greater extent during an initial challenge in FHP than FHN subjects. A required substrate for catalase dependent alcohol metabolism, hydrogen peroxide, is derived from fatty acid oxidation. The addition of oleate either in vivo or to the perfused liver can significantly stimulate catalase dependent alcohol metabolism by increasing hydrogen peroxide production (Bradford et al., 1999) (Handler & Thurman, 1997). Kaphailia demonstrated that fatty acid ethyl esters (FAEE) were elevated significantly after an acute exposure to ethanol as compared to a control group in which no FAEEs were detected (Kaphalia et al., 2004). Furthermore, the appearance of 16:0, 18:0 and 18:1 fatty acids in plasma support the hypothesis that an increase in plasma fatty acid supply is present in human subjects after acute ethanol. Rates of production of malondialdehyde (MDA), a product of lipid peroxidation, in erthryocytes of male patients who were either FHN or FHP for alcoholism were examined (Bidder & Jaeger, 1982). This study assessed patients 10 days after admission to a treatment center and thus the subjects were not intoxicated. The results from this study indicate that subjects who are FHP have twice the level of MDA produced by erthyrocytes when exposed to hydrogen peroxide. Thus, the findings from these two studies suggest that those subjects who are FHP may be more likely to have elevated hydrogen peroxide levels after an acute dose of alcohol than FHN subjects and would have elevated rates of catalase dependent alcohol metabolism. While it is not possible at this time to evaluate catalase, peroxide or fatty acid levels in the present study, there is significant evidence from work in rodents to suggest that rates of metabolism are regulated by these key components and more than one pathway of metabolism (Bradford et al., 1999;Bradford et al., 1993).

DIEM occurs in a subset of FHP subjects

Despite the differences in E_1 between FHP and FHN subjects, when statistically controlled for, FHP subjects were still more likely than FHN subjects to slow their rate of ethanol elimination. The observation that ethanol metabolism slows down in a significant portion of subjects at high-risk for alcoholism and is more likely to speed up in subjects at low-risk for alcoholism may have relevance for the development of tolerance to ethanol and, hence, to the risk of developing dependence. In animals, it has been shown that the duration of exposure to ethanol as well as the ascending rate of blood ethanol levels are associated with greater severity of physical dependence (Le Bourhis & Aufrere, 1983). We did not observe a difference between FHP and FHN subjects in time to peak ethanol level however, the extent of exposure to ethanol, as measured by area under the curve after Peak 2, was significantly greater in the subset of FHP-DIEM subjects (Table 2).

The increase in ethanol exposure observed in FHP-DIEM subjects is a function of two factorshigher peak BAC₂ and slower E_2 . The reason for the higher peak BAC₂ in FHP-DIEM subjects is not clear. However, it is noteworthy that FHP-DIEM subjects also had higher peak BAC₁ levels. Factors that increase peak ethanol levels after oral consumption include greater ethanol use, female gender, and an empty stomach. Our subjects were all male and had reported eating a light breakfast prior to the start of the experiment to minimize variation in initial gastric absorption. At the time of the second alcohol ingestion, all subjects had not eaten for about two hours, so food consumption should not have been a factor; however, cofactor supply for ADH-dependent or catalase dependent metabolism may have been limited. It is possible that FHP-DIEM subjects were heavier drinkers than FHN or FHP-NonDIEM subjects, despite similar self-reports of ethanol consumption, and that this explains their higher peak ethanol levels and their higher initial metabolic rate. It still does not account for their reduction in rate after the second alcohol consumption.

In humans, it has been a long-standing observation that rapid development of tolerance to ethanol is likely one factor in the development of alcohol dependence (Jellinek, 1960). Furthermore, the work of Schuckit (Schuckit, 1994) indicates that evidence of behavioral tolerance to ethanol early in life and prior to heavy drinking is a predictor of alcohol dependence later in life. Most likely, the major reason why some individuals develop tolerance to ethanol more rapidly than others is because of a differential neurobiologic response to ethanol. However, the duration and magnitude of exposure to ethanol is another factor that would be predicted to interact with neurobiologic adaptive systems to either hasten or slow the development of tolerance. Thus, one could hypothesize that FHP subjects with the DIEM phenomenon would be more likely to show behavioral tolerance to ethanol, in part, because of greater exposure to ethanol compared to FHN or other FHP subjects-- even when alcohol consumption is similar. The DIEM phenomenon would then represent another factor affecting the metabolism of ethanol with implications for risk of developing alcoholism--in addition to the evidence that genetic heterogeneity of alcohol dehydrogenase and aldehyde dehydrogenase affects risk for alcoholism (Higuchi et al., 1995).

Conclusions

Before the DIEM phenomenon can be accepted as a possible factor contributing to risk for alcoholism it will be necessary to replicate the core finding in high-risk and low-risk subjects with very careful assessment of prior consumption of ethanol. It will also be of interest to genotype subjects to examine for relationships between alcohol metabolizing pathways and the DIEM phenomenon. Examination of alcohol metabolism over the course of a continuous drinking period in DIEM vs. Non-DIEM subjects will also be of interest to more closely model what the effects of DIEM might be during more natural drinking behavior. Testing the hypothesis that DIEM is indeed associated with greater behavioral tolerance to ethanol will be necessary to determine if DIEM affects the behavioral response to ethanol.

In summary, we have described a phenomenon of a decrease in rate of ethanol metabolism following a priming dose of ethanol that is more prevalent in young men with an alcoholic father compared to young men without a family history of alcoholism. Though the reason for the DIEM phenomenon is unknown, DIEM could have implications for understanding the differential development of tolerance in high-risk vs. low-risk subjects and, thus, represent another factor in the complex path leading to alcoholism.

Acknowledgements

This work was completed with the assistance of grant MH33127 from the National Institute of Mental Health, Rockville, Md. and a grant from the North Carolina Alcoholism Research Authority. The authors would like to acknowledge and thank Dr. Dana Quade of the Biostatistics Department at the University of North Carolina, Chapel Hill for his excellent guidance on statistical evaluation of our data and the support of the nursing staff of the Clinical Research Unit, Dorothea Dix Hospital. The authors would like to dedicate this paper to the memory of the late Dr. Ronald G. Thurman, Ph.D, Professor of Pharmacology and Director of the Laboratory of Hepatobiology and Toxicology at the University of North Carolina, Chapel Hill.

Reference List

- Andreasen NC, Endicott J, Spitzer RL, Winokur G. The family history method using diagnostic criteria. Reliability and validity. Arch Gen Psychiatry 1977;34:1229-1235. [PubMed: 911222]
- Badger TM, Hoog JO, Svensson S, McGehee RE Jr, Fang C, Ronis MJ, Ingelman-Sundberg M. Cyclic expression of class I alcohol dehydrogenase in male rats treated with ethanol. Biochem Biophys Res Commun 2000;3:684–688. [PubMed: 10924336]
- Bidder TG, Jaeger PD. Malondialdehyde production by erythrocytes from alcoholic and non-alcoholic subjects. Life Sci 1982;30:1021-1027. [PubMed: 7070205]
- Bradford BU, Enomoto N, Ikejima K, Rose ML, Bojes HK, Forman DT, Thurman RG. Peroxisomes are involved in the swift increase in alcohol metabolism. J Pharmacol Exp Ther 1999;288:254-259. [PubMed: 9862778]
- Bradford BU, Seed CB, Handler JA, Forman DT, Thurman RG. Evidence that catalase is a major pathway of ethanol oxidation in vivo: dose-response studies in deer mice using methanol as a selective substrate. Arch Biochem Biophys 1993;303:172–176. [PubMed: 8489262]
- Choi IG, Kee BS, Son HG, Ham BJ, Yang BH, Kim SH, Lee JS, Son BK, Lee BY, Lee SY, Chai YG, Shin HD. Genetic polymorphisms of alcohol and aldehyde dehydrogenase, dopamine and seratonin transporters in familial and non-familial alcoholism. Eur Neuropsychopharmacol 2006;16:123-128. [PubMed: 16125912]
- Cloninger CR. Neurogenetic adaptive mechanisms in alcoholism. Science 1987;236:410-416. [PubMed: 2882604]
- Glassman EB, McLaughlin GA, Forman DT, Felder MR, Thurman RG. Role of alcohol dehydrogenase in the swift increase in alcohol metabolism (SIAM): studies with deermice deficient in alcohol dehydrogenase. Biochem Pharmacol 1989;34:3523-3526. [PubMed: 2932115]
- Handler JA, Bradford BU, Garbutt JC, Janowsky D, Miller L, Karnitschnig J, Seed CB, Forman DT, Thurman RG. Decrease in ethanol metabolism (DIEM) in sons of alcoholics and rodents. Alcohol Alcohol 1990;25:577-578.
- Handler, JA.; Thurman, RG. Catalase. In: Sipes, IG.; McQueen, CA.; Gandolfi, AJ., editors. Comprehensive Toxicology. 3. 1997. p. 217-227.
- Hasin DS, Grant BF. The co-occurrence of DSM-IV alcohol abuse in DSM-IV alcohol dependence: results of the National Epidemiologic Survey on Alcohol and Related Conditions on heterogeneity that differ by population subgroup. Arch Gen Psychiatry 2004;61:891–896. [PubMed: 15351767]
- Hesselbrock V, Begleiter H, Porjesz B, O'Connor S, Bauer L. P300 event-related potential amplitude as an endophenotype of alcoholism--evidence from the collaborative study on the genetics of alcoholism. J Biomed Sci 2001;8:77-82. [PubMed: 11173979]
- Higuchi S, Matsushita S, Murayama M, Takagi S, Hayashida M. Alcohol and aldehyde dehydrogenase polymorphisms and the risk for alcoholism. Am J Psychiatry 1995;152:1219-1221. [PubMed: 76254771
- Jellinek EM. Alcoholism, a genus and some of its species. Can Med Assoc J 1960;83:1341-1345. [PubMed: 13789799]
- Kaphalia BS, Cai P, Khan MF, Okorodudu AO, Ansari GA. Fatty acid ethyl esters: markers of alcohol abuse and alcoholism. Alcohol 2004;34:151-158. [PubMed: 15902908]
- Kendler KS, Heath AC, Neale MC, Kessler RC, Eaves LJ. A population-based twin study of alcoholism in women. JAMA 1992;268:1877-1882. [PubMed: 1404711]
- Koechling UM, Amit Z, Negrete JC. Family history of alcoholism and the mediation of alcohol intake by catalase: further evidence for catalase as a marker of the propensity to ingest alcohol. Alcohol Clin Exp Res 1995;19:1096-1104. [PubMed: 8561276]
- Lai CL, Chao YC, Chen YC, Liao CS, Chen MC, Liu YC, Yin SJ. No sex and age influence on the expression pattern and activities of human gastric alcohol and aldehyde dehydrogenases. Alcohol Clin Exp Res 2000;24:1625–1632. [PubMed: 11104109]
- Liangpunsakul S, Kolwankar D, Pinto A, Gorski JC, Hall SD, Chalasani N. Activity of CYP2E1 and CYP3A enzymes in adults with moderate alcohol consumption: a comparison with nonalcoholics. Hepatology 2005;41:1144-1150. [PubMed: 15841467]

- Le Bourhis B, Aufrere G. Pattern of alcohol administration and physical dependence. Alcohol Clin Exp Res 1983;7:378–381. [PubMed: 6362461]
- Levitt MD, Li R, DeMaster EG, Elson M, Furne J, Levitt DG. Use of measurements of ethanol absorption from stomach and intestine to assess human ethanol metabolism. Am J Physiol 1997;273:G951– G957. [PubMed: 9357840]
- Li T-K, Bosron WF, Dafeldecker WP, Lange LG, Vallee BL. Isolation of II-alcohol dehydrogenase of human liver: Is it a determinant of alcoholism? PNAS 1977;74:4378–4381. [PubMed: 270680]
- Ratsma JE, Van Der Stelt O, Gunning WB. Neurochemical markers of alcoholism vulnerability in humans. Alcohol Alcohol 2002;37:522–533. [PubMed: 12414542]
- Schuckit MA. Subjective responses to alcohol in sons of alcoholics and control subjects. Arch Gen Psychiat 1984;41:879–884. [PubMed: 6466047]
- Schuckit MA. Low level of response to alcohol as a predictor of future alcoholism. Am J Psychiatry 1994;151:184–189. [PubMed: 8296886]
- Schuckit MA, Tsuang JW, Anthenelli RM, Tipp JE, Nurnberger JI Jr. Alcohol challenges in young men from alcoholic pedigrees and control families: a report from the COGA project. J Stud Alcohol 1996;57:368–377. [PubMed: 8776678]
- Spitzer RL, Endicott J, Robins E. Research diagnostic criteria: rationale and reliability. Arch Gen Psychiatry 1978;35:773–782. [PubMed: 655775]
- Takahashi T, Lasker JM, Rosman AS, Lieber CS. Induction of cytochrome P-4502E1 in the human liver by ethanol is caused by a corresponding increase in encoding messenger RNA. Hepatology 1993;17:236–245. [PubMed: 8428720]
- Thurman, RG.; Bleyman, MA.; Yuki, T.; Bradford, BU. Genetic model to study the adaptive increase in ethanol metabolism due to prior exposure to ethanol in the rat. In: Eriksson, K.; Kilamaa, K., editors. Animal models in alcohol research. New York: Academic Press; 1980. p. 63-71.
- Thurman RG, Bradford BU, Glassman EB. The swift increase in alcohol metabolism (SIAM) in four inbred strains of mice. Pharm Biochem and Behavior 1983;18(Suppl 1):171–176.
- Thurman RG, Cheren I, Forman DT, Ewing JA, Glassman EB. Swift increase in alcohol metabolism in humans. Alcohol Clin Exp Res 1989;13:572–576. [PubMed: 2679213]
- Vieten C, Seaton KL, Feiler HS, Wilhelmsen KC. The University of California, San Francisco Family Alcoholism Study. I. Design, methods, and demographics. Alcohol Clin Exp Res 2004;28:1509– 1516. [PubMed: 15597083]
- Widmark, EP. Die Theoretischen Grundlagen und die praktische verwendbarkeit der gerichtmedizinische alkoholbestimmung. Berlin: Urban and Schwarzenberg; 1932.
- Wilson JR, Erwin VG, McClearn GE. Effects of ethanol: I. Acute metabolic tolerance and ethnic differences. Alcohol Cin Exp Res 1984;8:226–232.
- Yuki T, Bradford BU, Thurman RG. Role of hormones in the mechanism of the swift increase in alcohol metabolism in the rat. Pharmacol Biochem Behav 1980;13:67–71. [PubMed: 6113604]
- Yuki T, Thurman RG. The swift increase in alcohol metabolism: Time course for the increase in hepatic oxygen uptake and the involvement of glycolysis. Biochem J 1980;186:119–126. [PubMed: 6989357]

NIH-PA Author Manuscript

Bradford et al.



Figure 1.

Typical experiment in a Family History Positive Subject showing the two ascending ethanol phases, the two peak BAC levels and the two descending ethanol phases. The first rate of ethanol metabolism is calculated at 141 mg/kg/hr and the second rate at 94 mg/kg/hr with the change in rate being a reduction of 33%.





Bradford et al.

NIH-PA Author Manuscript





Bradford et al.



Figure 4.

Dot Plot of Area Under Curve following Second Peak Ethanol as an Estimate of Ethanol Exposure in FHP-DIEM (n=12), FHP-Non DIEM (n=11), and FHN (n=15) Subjects.

Table 1 Comparison of FHN and FHP Subjects on Demographic and Ethanol Metabolism Measures

Measure	FHN (n=15) Mean ± S.D.	FHN (n=23) Mean ± S.D.	t-test p-value
Age (yrs.)	22 ± 1	23 ± 3	.19
Height/Weight Ratio (cm/kg)	1.9 ± 0.1	1.9 ± 0.2	.74
Alcohol Use (Drinks/30 d)	38 ± 21	33 ± 23	.50
Nicotine Use (Cigarettes/30 d)	0 ± 0	7 ± 25	.18
Lifetime Marijuana Use	82 ± 258	60 ± 135	.76
Peak BAC ₁ (mg/dl)	82 ± 13	91 ± 16	.06
$E_1 (mg/kg/hr)$	124 ± 18	148 ± 36	.01
Peak BAC ₂ (mg/dl)	88 ± 15	97 ± 19	.13
$E_2 (mg/kg/hr)$	134 ± 30	121 ± 29	.19
Percent Rate Change	$+10 \pm 27$	-15 ± 24	.007
AUC (mg/dl min)	$14,676 \pm 5344$	$16,628 \pm 5993$.30

Table 2

Comparison of FHN, FHP-DIEM and FHP-NonDIEM Subjects on Demographic and Ethanol Metabolism Measures

Measure	FHN(n=15) Mean ± S.D.	FHP-DIEM(n=12) Mean ± S.D.	FHP-NonDIEM (n=11) Mean ± S.D.	F-test p-value
Age (yrs.)	22 ± 1	24 ± 4	21 ± 2	.02
Ht/Wt Ratio (cm/kg)	1.9 ± 0.1	1.9 ± 0.2	1.9 ± 0.1	.72
Alcohol Use (Drinks/30 d)	38 ± 21	34 ± 21	32 ± 27	.80
Peak BAC ₁ (mg/dl)	82 ± 13	99 ± 14	83 ± 13	.005
$E_1 (mg/kg/hr)$	124 ± 18	161 ± 41	133 ± 25	.008
Peak BAC ₂ (mg/dl)	88 ± 15	105 ± 18	89 ± 16	.03
$E_2 (mg/kg/hr)$	134 ± 30	107 ± 31	137 ± 18	.02
Percent Rate Change	10 ± 27	-33 ± 13	4 ± 16	.0001
AUC (mg/dl min)	$14,676 \pm 5344$	$19,792 \pm 6003$	$13,177 \pm 3778$.01