

References

Edmunds, J., Miles, S. C., & Fulbrook, P. (2011). Tongue-tie and breastfeeding: a review of the literature. *Breastfeeding review: professional publication of the Nursing Mothers' Association of Australia*, 19(1), 19–26.

Segal, L. M., Stephenson, R., Dawes, M., & Feldman, P. (2007). Prevalence, diagnosis, and treatment of ankyloglossia: methodologic review. *Canadian Family Physician*, 53(6), 1027–1033.

23 Articles in Systematic Review

Bellinger, V., Solari, D., Hogan, M., Rodda, K., Shadbolt, B., & Todd, D. (2018). Tongue-tie division in the newborn: Follow-up at 9 and 38 months. *Breastfeeding Review*, 26(1), 13–22.

Berry, J., Griffiths, M., & Westcott, C. (2012). A double-blind, randomized, controlled trial of tongue-tie division and its immediate effect on breastfeeding. *Breastfeeding Medicine : The Official Journal of the Academy of Breastfeeding Medicine*, 7(3), 189–193.
<https://doi.org/10.1089/bfm.2011.0030>

Billington, J., Yardley, I., & Upadhyaya, M. (2018). Long-term efficacy of a tongue tie service in improving breast feeding rates: A prospective study. *Journal of Pediatric Surgery*, 53(2), 286–288. <https://doi.org/10.1016/j.jpedsurg.2017.11.014>

Braccio, S., Chadderton, Z., Sherridan, A., & Upadhyaya, M. (2016). Tongue-tie division. Is it worth it? A retrospective cohort study. *British Journal of Midwifery*, 24(5), 317–321.
<https://doi.org/10.12968/bjom.2016.24.5.317>

Buryk, M., Bloom, D., & Shope, T. (2011). Efficacy of neonatal release of ankyloglossia: a randomized trial. *Pediatrics*, 128(2), 280–288. <https://doi.org/10.1542/peds.2011-0077>

Dollberg, S., Marom, R., & Botzer, E. (2014). Lingual frenotomy for breastfeeding difficulties: a prospective follow-up study. *Breastfeeding Medicine : The Official Journal of the Academy of Breastfeeding Medicine*, 9(6), 286–289. <https://doi.org/10.1089/bfm.2014.0010>

Emond, A., Ingram, J., Johnson, D., Blair, P., Whitelaw, A., Copeland, M., & Sutcliffe, A. (2014). Randomised controlled trial of early frenotomy in breastfed infants with mild-moderate tongue-tie. *Archives of Disease in Childhood. Fetal and Neonatal Edition*, 99(3), F189-95.
<https://doi.org/10.1136/archdischild-2013-305031>

Illing, S., Minnee, M., Wheeler, J., & Illing, L. (2019). The value of frenotomy for ankyloglossia from a parental perspective. *The New Zealand Medical Journal*, 132(1500), 70–81.

Martinelli, R. L. de C., Marchesan, I. Q., Gusmão, R. J., Honório, H. M., & Berretin-Felix, G. (2015). The effects of frenotomy on breastfeeding. *Journal of Applied Oral Science : Revista FOB*, 23(2), 153–157. <https://doi.org/10.1590/1678-775720140339>

McGoldrick, R., Solari, D., Hogan, M., Corrigan, I., Cutting, A., Shadbolt, B., & Todd, D. A. (2016). Tongue-tie in the newborn: Follow-up in the first 6 months. *Breastfeeding Review : Professional Publication of the Nursing Mothers' Association of Australia*, 24(3), 33–40.

Mettias, B., O'Brien, R., Abo Khatwa, M. M., Nasrallah, L., & Doddi, M. (2013). Division of tongue tie as an outpatient procedure. Technique, efficacy and safety. *International Journal of Pediatric Otorhinolaryngology*, 77(4), 550–552. <https://doi.org/10.1016/j.ijporl.2013.01.003>

Miller, A. S., & Miller, J. E. (2017). Is tongue tie really the problem? Incidence of ankyloglossia in an infant population presented with suboptimal feeding: a cross-sectional survey. *Journal of Clinical Chiropractic Pediatrics*, 16(1), 1350–1354.

Miranda, B. H., & Milroy, C. J. (2010). A quick snip - A study of the impact of outpatient tongue tie release on neonatal growth and breastfeeding. *Journal of Plastic, Reconstructive & Aesthetic Surgery*, 63(9), e683-5. <https://doi.org/10.1016/j.bjps.2010.04.003>

Muldoon, K., Gallagher, L., McGuinness, D., & Smith, V. (2017). Effect of frenotomy on breastfeeding variables in infants with ankyloglossia (tongue-tie): a prospective before and after cohort study. *BMC Pregnancy and Childbirth*, 17(1), 373. <https://doi.org/10.1186/s12884-017-1561-8>

O'Callahan, C., Macary, S., & Clemente, S. (2013). The effects of office-based frenotomy for anterior and posterior ankyloglossia on breastfeeding. *International Journal of Pediatric Otorhinolaryngology*, 77(5), 827–832. <https://doi.org/10.1016/j.ijporl.2013.02.022>

Page, T. A., Lai, J., Zheng, L., Webster, K. L., & Gill, A. C. (2019). Retrospective quality audit of frenotomy for neonatal tongue-tie. *Breastfeeding Review*, 27(1), 13–19.

Ramoser, G., Guóth-Gumberger, M., Baumgartner-Sigl, S., Zoeggeler, T., Scholl-Bürgi, S., & Karall, D. (2019). Frenotomy for tongue-tie (frenulum linguae breve) showed improved symptoms in the short- and long-term follow-up. *Acta Paediatrica*, 108(10), 1861–1866. <https://doi.org/10.1111/apa.14811>

Schlatter, S.-M., Schupp, W., Otten, J.-E., Harnisch, S., Kunze, M., Stavropoulou, D., & Hentschel, R. (2019). The role of tongue-tie in breastfeeding problems-A prospective observational study. *Acta Paediatrica*, 108(12), 2214–2221. <https://doi.org/10.1111/apa.14924>

Sethi, N., Smith, D., Korteque, S., Ward, V. M. M., & Clarke, S. (2013). Benefits of frenulotomy in infants with ankyloglossia. *International Journal of Pediatric Otorhinolaryngology*, 77(5), 762–765. <https://doi.org/10.1016/j.ijporl.2013.02.005>

Sharma, S. D., & Jayaraj, S. (2015). Tongue-tie division to treat breastfeeding difficulties: our experience. *The Journal of Laryngology and Otology*, 129(10), 986–989. <https://doi.org/10.1017/S002221511500225X>

Srinivasan, A., Al Khoury, A., Puzhko, S., Dobrich, C., Stern, M., Mitnick, H., & Goldfarb, L. (2019). Frenotomy in Infants with Tongue-Tie and Breastfeeding Problems. *Journal of Human Lactation*, 35(4), 706–712. <https://doi.org/10.1177/0890334418816973>

Steehler, M. W., Steehler, M. K., & Harley, E. H. (2012). A retrospective review of frenotomy in neonates and infants with feeding difficulties. *International Journal of Pediatric Otorhinolaryngology*, 76(9), 1236–1240. <https://doi.org/10.1016/j.ijporl.2012.05.009>

Wakhanrittee, J., Khorana, J., & Kiatipunsodsai, S. (2016). The outcomes of a frenulotomy on breastfeeding infants followed up for 3 months at Thammasat University Hospital. *Pediatric Surgery International*, 32(10), 945–952. <https://doi.org/10.1007/s00383-016-3952-8>