This is an Accepted Manuscript for *Infection Control & Hospital Epidemiology* as part of the Cambridge Coronavirus Collection. DOI: 10.1017/ice.2020.1427

## **Article Type: Commentary**

**Title:** Lessons Learned in Infection Prevention for Ebola Virus Disease and the COVID-19 Pandemic—Principles Underlying Prevention

## Running title: Principles Underlying Prevention

Erica S. Shenoy, MD, PhD<sup>1, 2, 3</sup>

David J. Weber, MD, MPH<sup>4,5</sup>

<sup>1</sup> Infection Control Unit, Massachusetts General Hospital, Boston, MA

<sup>2</sup> Division of Infectious Diseases, Massachusetts General Hospital, Boston, MA

<sup>3</sup> Harvard Medical School, Boston, MA

<sup>4</sup> Division of Infectious Disease, School of Medicine, University of North Carolina at Chapel

Hill, Chapel Hill, NC, 27599-7030

<sup>5</sup> Department of Hospital Epidemiology, UNC Medical Center, Chapel Hill, NC, 27154

Keywords: Ebola virus disease, SARS-CoV-2, healthcare personnel infection, prevention

Word count: 1143/1200

Figure: 1

Table: 1

<sup>†</sup> Corresponding author: Erica S. Shenoy, MD, PhD Massachusetts General Hospital 55 Fruit St., Gr-J 504 Boston, MA 02114 Phone: 617-643-5637 Fax: 617-724-0267 eshenoy@mgh.harvard.edu Delivery of care to patients with highly communicable diseases balances the potential risk of transmission from the patient-to-healthcare personnel (HCP) with the risks to the patient of delayed or reduced access to needed interventions. The risk of transmission to HCP depends on many factors, described by the chain of transmission (**Figure**), and include the establishment of a reservoir (human, animal, inanimate environment), exit of the infectious agent from the reservoir and survival in the environment, with transmission by direct or indirect contact, droplet, airborne modes or combinations of these modes, and finally entry of the infectious agent via a portal of entry to a susceptible host at an inoculum sufficient to establish infection. Efforts to prevent transmission in healthcare settings—between patients, visitors, and HCP—are all aimed at interrupting the chain of transmission and include, in addition to correct and consistent use of personal protective equipment (PPE) plus rapid institution of appropriate isolation precautions as indicated by the mode of transmission, multiple other interventions that minimize the risk of nosocomial transmission, often framed as part of the Hierarchy of Controls applied to HCP safety, but also with applications for reducing overall risk of transmission to patients and visitors.<sup>1</sup>

In this issue of *Infection Control and Hospital Epidemiology*, DiLorenzo et al report on a survey of policies of Ebola Treatment Centers (ETCs) with respect to provision (planned or actual) of critical care interventions for patients with Viral Hemorrhagic Fevers (VHF) such as Ebola Virus Disease (EVD).<sup>2</sup> The authors distributed a 58-item survey to 82 ETCs between January 2020-March 2020, and report on the responses of 17 institutions of which fewer than half had experience caring for patients with VHFs or persons under investigation (PUIs). The authors queried institutions on policies in nine critical care areas (renal replacement therapy, endotracheal intubation and mechanical ventilation, extracorporeal membrane oxygenation, chest compressions, pharmacological cardioversion, electrical cardioversion, defibrillation, cricothyrorotomy, and code status) as well as to what extent staff safety, lack of appropriate technology, lack of clinical guidelines, clinical futility, and limitations of the environment of care, influenced policies regarding provision of care. A majority of respondents had policies with respect to renal replacement, endotracheal intubutation and mechanical ventilation, and chest compressions, although applications of each varied by patient level factors. For other interventions, fewer respondents reported having policies, and amongst those there was variation in types of patients (PUI vs confirmed VHF) to whom it would be offered. Among the factors

influencing decision regarding offering care to either PUIs or confirmed VHF patients, staff safety and clinical futility were reported to impact decisions "somewhat" or "greatly" for a majority of respondents whereas lack of appropriate technology, guidelines, or physical limitations in the environment of care either did not limit care or limited minimally for the majority of respondents.

The fact that healthcare personnel safety has such a prominent impact on decisions to offer particular types of care is not unique to VHFs, and has been a concern raised in the provision of care to patients infected with SARS-CoV-2, there are some prominent differences between the two pathogens that underly the potential risk of nosocomial acquisition to HCP. Specifically, the primary modes of transmission (EVD primarily contact and SARS-CoV-2 primarily droplet), and the ability of individuals to transmit infection while asymptomatic or presymptomatic (not considered likely with EVD, prominent with SARS-CoV-2) are key aspects that inform the infection prevention strategies (**Table**).

The available data from VHFs and SARS-CoV-2, however, demonstrate that the major risks to HCP are from failure to identify patients at entrance to a healthcare facility as possibly infected and isolate them appropriately, failure to utilize personal protective equipment (PPE) correctly especially during donning and doffing, and inadequate PPE due to shortages. These same challenges have been present during the COVID-19 pandemic. Additionally, HCP-to-HCP spread of SARS-CoV-2 has been linked to lapses in masking and distancing where masks are removed to eat or drink such as breakrooms and nursing stations and physical distancing is not maintained. Acquistion by HCP has then led in some cases to transmission to patients with propagation of transmission.<sup>3</sup>

The primary intervention to reduce risk of nosocomial transmission relies on early identification of PUIs and initiation of isolation. For both EVD<sup>4</sup> and SARS-CoV-2<sup>5-8</sup>, failures at this critical juncture have resulted in exposures to HCP and transmission events. Failure to use appropriate PPE, closely tied to early identification of patients as PUIs and then use of the correct PPE. While self- and cross- contamination is a concern with SARS-CoV-2, and careful doffing and use of hand hygiene must be emphasized, transmission directly attributable to doffing failures has not been documented. One SARS-CoV-2 serological study failed to identify an association between positive serology and care of patients with COVID-19, however did note a strong association of living in a household with an individuals with suspected or confirmed SARS-

CoV-2 infection.<sup>9</sup> Another serological study of HCP noted lower prevalence of SARS-CoV-2 antibodies among HCP who reported consistent use of a facemask when caring for patients.<sup>10</sup> In contrast, due to contact with blood and body fluids as the primary mode of transmission with EVD, self- and cross-contamination is a priority concern to the extent that extensive training in the use of PPE, careful selection of PPE components and order of doffing, close attention to the design of the physical space where doffing occurs is warranted, and the implementation of a Trained Observer is recommended by the CDC to ensure each HCP doffs correctly and that instances of possible contamination are identified during the process and mitigated.<sup>11-15</sup> use of dedicated HCP with who have trained and exercised in the use of PPE for Ebola is recommended both to the high-risk aspect of doffing PPE while avoiding self- and cross-contamination, as well as the fact that the PPE used for EVD and other viral hemorrhagic fevers is not used routinely in most healthcare settings. Adjunctive approaches, such as techniques to visualize contamination<sup>16</sup> and the use of ultraviolet disinfection of  $PPE^{17}$  have been assessed to reduce the risk to HCP. In some settings with EVD, and world-wide with SARS-CoV-2 due to the large-scale global nature of the pandemic with resultant interruption of supply chain, PPE shortages have led to strategies that have included extended use, re-use of PPE following disinfection, and use of alternative PPE components that have not been certified, as well as lack of adequate PPE.<sup>18</sup> HCP-to-HCP transmission of VHFs has been reported.<sup>19,20</sup> HCP-to-HCP of SARS-CoV-2 infection has been well documented via droplet spread, in part, because it can be transmitted from asymptomatic, pre-symptomatic, and pauci-symptomatic individualsespecially in settings where masking is not present such as breakrooms.<sup>21,22</sup> Transmission events are not restricted to HCP interactions in the workplace are more likely to occur during external activities such as commuting while unmasked and other social activities where masking compliance between HCP may be reduced. DiLorenzo et al demonstrate that HCP safety in provision of critical care to EVD PUIs is informed by assessment of risk of potential for patient-to-HCP transmission. Similar concerns have underscored the COVID-19 pandemic, and highlight the importance of multi-faceted approaches to interrupting the chain of transmission. Differences between the two pathogens, however, specifically the primary modes of transmission and role of asymptomatic/presymptomatic transmission underscore differences observed in the overall risk of patient-to-HCP transmission.

Figure. The Chain of Transmission.

Figure Legend. Transmission from one individual to another requires completion of each step in the chain of transmission. Beginning with an infectious agent in a reservoir (e.g., human, animal, or inanimate object/surface), the infectious agent must exit the reservoir through portal of exit, survive in the environment, and be transmitted by contact, droplet, or airborne routes (or a combination thereof), enter as susceptible host through a portal of entry (e.g., eyes, nose, mouth, wound) at an inoculum sufficient to establish infection.



<b>Table.</b> Comparison of Pathogens Primarily Transmitted by Contact with Body Fluids (e.g., Ebola virus)			
Versus Respiratory Droplets and Droplet Nuclei (e.g., SARS-CoV-2)			
	Ebola virus	SARS-CoV-2	
Microbiology			
Year identified	1976	2019	
Family	Filaviridae	Coronaviridae	
Genome	RNA	RNA	
Coat	Enveloped	Enveloped	
Epidemiology			
Prevalence	Repeated outbreaks	Pandemic	
Reservoir	Bats	Bats; research ongoing to identify	
		additional potential reservoirs	
Intermediate host	Primates and other animals	None demonstrated	
Primary mode of	Direct Contact: Contact with	Respiratory droplets and short range	
transmission	infectious body fluids	droplet nuclei	
Other modes of	Indirect contact (i.e., contaminated	Direct and indirect contact (i.e.,	
transmission	surfaces, devices), sexual, blood	contaminated surfaces, devices)	
	transfusion		
Basic reproductive rate	1.5-2.0 <sup>23</sup>	1.8-3.6 <sup>24</sup>	
(R <sub>0</sub> )			
Asymptomatic and	No	Yes	
presymptomatic			
transmission			
Incubation period	6-12 days (range, 2-21)	2-14 days	
Case-fatality rate	~50% (range, 25%-90%)	~15% among hospitalized patients	
Treatment	Monoclonal antibody combination	Remdesivir; bamlanivimab	
	(atoltivimab, maftivimab, and		
	odesivimab-ebgn)		
Infection Prevention			
Nosocomial transmission	Yes	Yes	

involving HCP (HCP-to-		
HCP, HCP-to-patient,		
patient-to-HCP)		
Laboratory biosafety level	BSL-4	BSL-3
Survival on surfaces	Hours to a few day	Hours to a few days
Antiseptic	60%-90% alcohol based-product	60%-90% alcohol based-product
Disinfectant	EPA, emerging virus claim (List "N")	EPA, emerging virus claim (List "N")
Special handling of used linens, patient waste	Yes	No
PPE worn by HCP (CDC)	<ol> <li>Single-use (disposable) fluid- resistant gown that extends to at least mid-calf or single-use (disposable) fluid-resistant coveralls without integrated hood</li> <li>Single-use (disposable) full face shield</li> <li>Single-use (disposable) facemask</li> <li>Single-use (disposable) gloves with extended cuffs. Two pairs of gloves should be worn. At a minimum, outer gloves should have extended cuffs.<sup>25</sup></li> </ol>	<ol> <li>N95 respirator (or equivalent or higher-level respirator) or facemask (if a respirator is not available)</li> <li>Eye protection (i.e., goggles or a face shield that covers the front and sides of the face)</li> <li>Single use (disposable), clean, non-sterile gloves</li> <li>Single use (disposable) isolation gown or cloth gown.<sup>26</sup></li> </ol>
Pre-exposure prophylaxis	Vaccine	None
Post-exposure prophylaxis	None approved for post-exposure prophylaxis	None

BSL, biosafety level; EPA, U.S. Environmental Protection Agency; HCP, healthcare personnel;

## REFERENCES

- NIOSH. Hierarchy of Controls. <u>https://www.cdc.gov/niosh/topics/hierarchy/default.html</u>. Published 2015. Accessed.
- DiLorenzo MA BC, Herstein JJ, Evans L, Lowe JJ, Gibbs SG, Bhadelia N. Institutional Policies and Readiness in Management of Critical Illness in Among Patients with Viral Hemorrhagic Fever. *Infect Control Hospital Epidemiol.* 2020.
- CDC. Healthcare Facilities: Managing Operations During the COVID-19 Pandemic. <u>https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-hcf.html</u>. Published 2020. Accessed.
- Anderson-Fletcher E, Vera D, Abbott J. The Texas Health Presbyterian Hospital Ebola Crisis: A Perfect Storm of Human Errors, System Failures, and Lack of Mindfulness. In:2015.
- Bays DJ, Nguyen MH, Cohen SH, et al. Investigation of Nosocomial SARS-CoV-2 Transmission from Two Patients to Health Care Workers Identifies Close Contact but not Airborne Transmission Events. *Infection control and hospital epidemiology*. 2020:1-22.
- 6. Baker MA, Rhee C, Fiumara K, et al. COVID-19 infections among HCWs exposed to a patient with a delayed diagnosis of COVID-19. *Infection control and hospital epidemiology*. 2020;41(9):1075-1076.
- Heinzerling A, Stuckey MJ, Scheuer T, et al. Transmission of COVID-19 to Health Care Personnel During Exposures to a Hospitalized Patient - Solano County, California, February 2020. MMWR Morb Mortal Wkly Rep. 2020;69(15):472-476.
- 8. Ng K, Poon BH, Kiat Puar TH, et al. COVID-19 and the Risk to Health Care Workers: A Case Report. *Annals of internal medicine*. 2020;172(11):766-767.
- Steensels D, Oris E, Coninx L, et al. Hospital-Wide SARS-CoV-2 Antibody Screening in 3056 Staff in a Tertiary Center in Belgium. *Jama*. 2020;324(2):195-197.
- Self WH, Tenforde MW, Stubblefield WB, et al. Seroprevalence of SARS-CoV-2 Among Frontline Health Care Personnel in a Multistate Hospital Network - 13 Academic Medical Centers, April-June 2020. MMWR Morb Mortal Wkly Rep. 2020;69(35):1221-1226.
- CDC. Guidance on Personal Protective Equipment (PPE) To Be Used By Healthcare Workers during Management of Patients with Confirmed Ebola or Persons under

Investigation (PUIs) for Ebola who are Clinically Unstable or Have Bleeding, Vomiting, or Diarrhea in U.S. Hospitals, Including Procedures for Donning and Doffing PPE. 2015.

- Rock C, Cosgrove SE, Keller SC, et al. Using a Human Factors Engineering Approach to Improve Patient Room Cleaning and Disinfection. *Infection control and hospital epidemiology*. 2016;37(12):1502-1506.
- Reddy SC, Valderrama AL, Kuhar DT. Improving the Use of Personal Protective Equipment: Applying Lessons Learned. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2019;69(Suppl 3):S165-S170.
- 14. DuBose JR, Matić Z, Sala MFW, et al. Design strategies to improve healthcare worker safety in biocontainment units: learning from ebola preparedness. *Infection control and hospital epidemiology*. 2018;39(8):961-967.
- Wong MF, Matić Z, Campiglia GC, et al. Design Strategies for Biocontainment Units to Reduce Risk During Doffing of High-level Personal Protective Equipment. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2019;69(Suppl 3):S241-S247.
- Tomas ME, Cadnum JL, Mana TS, et al. Utility of a Novel Reflective Marker Visualized by Flash Photography for Assessment of Personnel Contamination During Removal of Personal Protective Equipment. *Infection control and hospital epidemiology*. 2016;37(6):711-713.
- Tomas ME, Cadnum JL, Jencson A, Donskey CJ. The Ebola disinfection booth: evaluation of an enclosed ultraviolet light booth for disinfection of contaminated personal protective equipment prior to removal. *Infection control and hospital epidemiology*. 2015;36(10):1226-1228.
- CDC. Strategies for Optimizing Personal Protective Equipment (PPE) Supplies. <u>https://www.cdc.gov/coronavirus/2019-ncov/hcp/ppe-strategy/index.html</u>. Published 2020. Accessed.
- Burney MI, Ghafoor A, Saleen M, Webb PA, Casals J. Nosocomial outbreak of viral hemorrhagic fever caused by Crimean Hemorrhagic fever-Congo virus in Pakistan, January 1976. *The American journal of tropical medicine and hygiene*. 1980;29(5):941-947.

- 20. Ftika L, Maltezou HC. Viral haemorrhagic fevers in healthcare settings. *The Journal of hospital infection*. 2013;83(3):185-192.
- 21. Çelebi G, Pişkin N, Çelik Bekleviç A, et al. Specific risk factors for SARS-CoV-2 transmission among health care workers in a university hospital. *American journal of infection control.* 2020;48(10):1225-1230.
- 22. Ellsworth M, Chang M, Ostrosky-Zeichner L. Mind the gap: The hospital breakroom. *American journal of infection control.* 2020;48(10):1285.
- 23. Khan A, Naveed M, Dur-E-Ahmad M, Imran M. Estimating the basic reproductive ratio for the Ebola outbreak in Liberia and Sierra Leone. *Infect Dis Poverty*. 2015;4:13.
- 24. Petersen E, Koopmans M, Go U, et al. Comparing SARS-CoV-2 with SARS-CoV and influenza pandemics. *The Lancet Infectious diseases*. 2020;20(9):e238-e244.
- 25. CDC. For US Healthcare Settings: Donning and Doffing Personal Protective Equipment (PPE) for Evaluating Persons Under Investigation PUIs) for Ebola Who Are Clinically Stable and Do Not Have Bleeding, Vomiting, or Diarrhea. <u>https://www.cdc.gov/vhf/ebola/healthcare-us/ppe/guidance-clinically-stable-puis.html</u>. Published 2018. Accessed.
- 26. CDC. Interim Infection Prevention and Control Recommendations for Healthcare Personnel During the Coronavirus Disease 2019 (COVID-19) Pandemic. <u>https://www.cdc.gov/coronavirus/2019-ncov/hcp/infection-control-recommendations.html</u>. Published 2020. Accessed.